
Topology of surface singularities: superisolated, Lê-Yomdin and weighted Lê-Yomdin

Enrique ARTAL BARTOLO

Departamento de Matemáticas
Facultad de Ciencias
Instituto Universitario de Matemáticas y sus Aplicaciones
Universidad de Zaragoza

GKŁW Seminar in Singularity Theory
March 24th 2023

Funded by MCIN/AEI/10.13039/501100011033 (PID2020-114750GB-C31) and Gobierno de Aragón (E22_20R: "Álgebra y Geometría").



Surfaces in $\mathbb{C}^3$

Superisolated and Lê-Yomdin singularities

Zariski tuples and Jordan form

Zariski tuples of non-Alexander type

Quasi-homogeneous and Weighted Lê-Yomdin singularities



Abstract topology

- Ground field $\mathbb{C}$.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}_\varepsilon^{2N-1} \pitchfork S$.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}^{2N-1}_{\varepsilon} \pitchfork S$.
- *Link* of $(S,0)$: the topology of $K \equiv K_\varepsilon := \mathbb{S}^{2N-1}_{\varepsilon} \cap S$ does not depend on $\varepsilon \in (0,\varepsilon_0]$.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}^{2N-1}_{\varepsilon} \pitchfork S$.
- *Link* of $(S,0)$: the topology of $K \equiv K_\varepsilon := \mathbb{S}^{2N-1}_{\varepsilon} \cap S$ does not depend on $\varepsilon \in (0,\varepsilon_0]$.
- K is a 3-manifold, a graph manifold, S is topologically a cone over K.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}^{2N-1}_{\varepsilon} \pitchfork S$.
- *Link* of $(S,0)$: the topology of $K \equiv K_\varepsilon := \mathbb{S}^{2N-1}_{\varepsilon} \cap S$ does not depend on $\varepsilon \in (0,\varepsilon_0]$.
- K is a 3-manifold, a graph manifold, S is topologically a cone over K.
- A resolution of $(S,0)$ is a proper birational map $\sigma : (X,E) \to (S,0)$, X smooth, $\sigma_| : X \setminus E \overset{\cong}{\to} S \setminus \{0\}$.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}^{2N-1}_{\varepsilon} \pitchfork S$.
- *Link* of $(S,0)$: the topology of $K \equiv K_\varepsilon := \mathbb{S}^{2N-1}_{\varepsilon} \cap S$ does not depend on $\varepsilon \in (0,\varepsilon_0]$.
- K is a 3-manifold, a graph manifold, S is topologically a cone over K.
- A resolution of $(S,0)$ is a proper birational map $\sigma : (X,E) \to (S,0)$, X smooth, $\sigma_| : X \setminus E \overset{\cong}{\to} S \setminus \{0\}$.
- A resolution is *good* if $\sigma^{-1}(0)$ is a normal crossing divisor (smooth irreducible components and locally monomial).



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}^{2N-1}_\varepsilon \pitchfork S$.
- *Link* of $(S,0)$: the topology of $K \equiv K_\varepsilon := \mathbb{S}^{2N-1}_\varepsilon \cap S$ does not depend on $\varepsilon \in (0,\varepsilon_0]$.
- K is a 3-manifold, a graph manifold, S is topologically a cone over K.
- A resolution of $(S,0)$ is a proper birational map $\sigma : (X,E) \to (S,0)$, X smooth, $\sigma_| : X \setminus E \overset{\cong}{\to} S \setminus \{0\}$.
- A resolution is *good* if $\sigma^{-1}(0)$ is a normal crossing divisor (smooth irreducible components and locally monomial).
- Γ dual graph, vertices weighted by genus and self-intersection.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}^{2N-1}_{\varepsilon} \pitchfork S$.
- *Link* of $(S,0)$: the topology of $K \equiv K_\varepsilon := \mathbb{S}^{2N-1}_{\varepsilon} \cap S$ does not depend on $\varepsilon \in (0,\varepsilon_0]$.
- K is a 3-manifold, a graph manifold, S is topologically a cone over K.
- A resolution of $(S,0)$ is a proper birational map $\sigma : (X,E) \to (S,0)$, X smooth, $\sigma_| : X \setminus E \overset{\cong}{\to} S \setminus \{0\}$.
- A resolution is *good* if $\sigma^{-1}(0)$ is a normal crossing divisor (smooth irreducible components and locally monomial).
- Γ dual graph, vertices weighted by genus and self-intersection.
- Γ is also the plumbing graph of K.



Abstract topology

- Ground field $\mathbb{C}$.
- $(S,0) \subset (\mathbb{C}^N,0)$ germ of isolated singularity of surface.
- $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0,\varepsilon_0]$, $\mathbb{S}^{2N-1}_{\varepsilon} \pitchfork S$.
- *Link* of $(S,0)$: the topology of $K \equiv K_\varepsilon := \mathbb{S}^{2N-1}_{\varepsilon} \cap S$ does not depend on $\varepsilon \in (0,\varepsilon_0]$.
- K is a 3-manifold, a graph manifold, S is topologically a cone over K.
- A resolution of $(S,0)$ is a proper birational map $\sigma : (X,E) \to (S,0)$, X smooth, $\sigma_| : X \setminus E \overset{\cong}{\to} S \setminus \{0\}$.
- A resolution is *good* if $\sigma^{-1}(0)$ is a normal crossing divisor (smooth irreducible components and locally monomial).
- Γ dual graph, vertices weighted by genus and self-intersection.
- Γ is also the plumbing graph of K.
- In general, it does not determine the analytic type.



An example


$$S = \{xyz + x^3 + y^3 + z^4 = 0\} \subset \mathbb{C}^3$$



An example

$S = \{xyz + x^3 + y^3 + z^4 = 0\} \subset \mathbb{C}^3$


- $\sigma_0 : (X_0, E_0) \to (S, 0)$ minimal (non *good*) resolution:




An example

$$S = \{xyz + x^3 + y^3 + z^4 = 0\} \subset \mathbb{C}^3$$

- $\sigma_0 : (X_0, E_0) \to (S, 0)$ minimal (non *good*) resolution:
- $\sigma_1 : (X_1, E_0 \cup E_1) \to (S, 0)$ minimal *good* resolution:







An example

$S = \{xyz + x^3 + y^3 + z^4 = 0\} \subset \mathbb{C}^3$

- $\sigma_0 : (X_0, E_0) \to (S, 0)$ minimal (non *good*) resolution:
- $\sigma_1 : (X_1, E_0 \cup E_1) \to (S, 0)$ minimal *good* resolution:

Plumbing (Neumann, Waldhausen)

- Vertex v weighted (g_v, e_v): M_v oriented $\mathbb{S}^1$-fiber bundle over a compact surface of genus g_v with Euler number e_v.
- Edge between v, w: empty two fiber solid tori in M_v, M_w and glue the boundaries interchanging meridians and fibers.
- *Normalized* graphs determine K topologically.
- Incidence matrix A_Γ; $\det S := \det(-A_\Gamma) \in \mathbb{Z}_{\geq 1}$.



An example

$S = \{xyz + x^3 + y^3 + z^4 = 0\} \subset \mathbb{C}^3$

- $\sigma_0 : (X_0, E_0) \to (S, 0)$ minimal (non *good*) resolution:
- $\sigma_1 : (X_1, E_0 \cup E_1) \to (S, 0)$ minimal *good* resolution:

Plumbing (Neumann, Waldhausen)

- Vertex v weighted (g_v, e_v): M_v oriented $\mathbb{S}^1$-fiber bundle over a compact surface of genus g_v with Euler number e_v.
- Edge between v, w: empty two fiber solid tori in M_v, M_w and glue the boundaries interchanging meridians and fibers.
- *Normalized* graphs determine K topologically.
- Incidence matrix A_Γ; $\det S := \det(-A_\Gamma) \in \mathbb{Z}_{\geq 1}$.



An example

$S = \{xyz + x^3 + y^3 + z^4 = 0\} \subset \mathbb{C}^3$

- $\sigma_0 : (X_0, E_0) \to (S, 0)$ minimal (non *good*) resolution:
- $\sigma_1 : (X_1, E_0 \cup E_1) \to (S, 0)$ minimal *good* resolution:

Plumbing (Neumann, Waldhausen)

- Vertex v weighted (g_v, e_v): M_v oriented $\mathbb{S}^1$-fiber bundle over a compact surface of genus g_v with Euler number e_v.
- Edge between v, w: empty two fiber solid tori in M_v, M_w and glue the boundaries interchanging meridians and fibers.
- *Normalized* graphs determine K topologically.
- Incidence matrix A_Γ; $\det S := \det(-A_\Gamma) \in \mathbb{Z}_{\geq 1}$.



An example

$S = \{xyz + x^3 + y^3 + z^4 = 0\} \subset \mathbb{C}^3$

- $\sigma_0 : (X_0, E_0) \to (S, 0)$ minimal (non *good*) resolution:
- $\sigma_1 : (X_1, E_0 \cup E_1) \to (S, 0)$ minimal *good* resolution:

Plumbing (Neumann, Waldhausen)

- Vertex v weighted (g_v, e_v): M_v oriented $\mathbb{S}^1$-fiber bundle over a compact surface of genus g_v with Euler number e_v.
- Edge between v, w: empty two fiber solid tori in M_v, M_w and glue the boundaries interchanging meridians and fibers.
- *Normalized* graphs determine K topologically.
- Incidence matrix A_Γ; $\det S := \det(-A_\Gamma) \in \mathbb{Z}_{\geq 1}$.



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity
- Milnor number $\mu := \dim_{\mathbb{C}} \mathbb{C}\{x, y, z\}/\operatorname{Jac} F < \infty$.



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity
- Milnor number $\mu := \dim_{\mathbb{C}} \mathbb{C}\{x, y, z\}/\operatorname{Jac} F < \infty$.
- Embedded topology: topological type of $(\mathbb{S}^5, K)$.



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity
- Milnor number $\mu := \dim_{\mathbb{C}} \mathbb{C}\{x, y, z\}/\operatorname{Jac} F < \infty$.
- Embedded topology: topological type of $(\mathbb{S}^5, K)$.
- Main tool, Milnor fibration $(0 < \eta \ll \varepsilon < \varepsilon_0)$:

$$\mathbb{S}^5 \setminus K \xrightarrow{\frac{f}{|f|}} \mathbb{S}^1 \qquad\qquad \overline{\mathbb{B}}^6_\varepsilon \cap f^{-1}(\mathbb{S}^1_\eta) \xrightarrow{f} \mathbb{S}^1_\eta$$



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity
- Milnor number $\mu := \dim_{\mathbb{C}} \mathbb{C}\{x, y, z\}/\operatorname{Jac} F < \infty$.
- Embedded topology: topological type of $(\mathbb{S}^5, K)$.
- Main tool, Milnor fibration $(0 < \eta \ll \varepsilon < \varepsilon_0)$:

$$\mathbb{S}^5 \setminus K \xrightarrow{\frac{f}{|f|}} \mathbb{S}^1 \qquad\qquad \overline{\mathbb{B}}^6_\varepsilon \cap f^{-1}(\mathbb{S}^1_\eta) \xrightarrow{f} \mathbb{S}^1_\eta$$

- Milnor fiber $F \simeq \bigvee_\mu \mathbb{S}^2$.



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity
- Milnor number $\mu := \dim_{\mathbb{C}} \mathbb{C}\{x, y, z\}/\operatorname{Jac} F < \infty$.
- Embedded topology: topological type of $(\mathbb{S}^5, K)$.
- Main tool, Milnor fibration $(0 < \eta \ll \varepsilon < \varepsilon_0)$:

$$\mathbb{S}^5 \setminus K \xrightarrow{\frac{f}{|f|}} \mathbb{S}^1 \qquad\qquad \overline{\mathbb{B}}^6_\varepsilon \cap f^{-1}(\mathbb{S}^1_\eta) \xrightarrow{f} \mathbb{S}^1_\eta$$

- Milnor fiber $F \simeq \bigvee_\mu \mathbb{S}^2$.
- Fibrations determined by $\rho : F \to F$, diffeomorphism well-defined up to isotopy (monodromy).



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity
- Milnor number $\mu := \dim_{\mathbb{C}} \mathbb{C}\{x, y, z\}/\operatorname{Jac} F < \infty$.
- Embedded topology: topological type of $(\mathbb{S}^5, K)$.
- Main tool, Milnor fibration $(0 < \eta \ll \varepsilon < \varepsilon_0)$:

$$\mathbb{S}^5 \setminus K \xrightarrow{\frac{f}{|f|}} \mathbb{S}^1 \qquad\qquad \overline{\mathbb{B}}^6_\varepsilon \cap f^{-1}(\mathbb{S}^1_\eta) \xrightarrow{f} \mathbb{S}^1_\eta$$

- Milnor fiber $F \simeq \bigvee_\mu \mathbb{S}^2$.
- Fibrations determined by $\rho : F \to F$, diffeomorphism well-defined up to isotopy (monodromy).
- Homological monodromy: $\rho^* : H^2(F; R) \to H^2(F; R)$, $R = \mathbb{Z}, \mathbb{Q}, \mathbb{C}$.



Embedded topology of hypersurfaces

- $N = 3$, $S = \{F = 0\}$, $F(x, y, z) \in \mathbb{C}\{x, y, z\}$ irreducible, isolated singularity
- Milnor number $\mu := \dim_{\mathbb{C}} \mathbb{C}\{x, y, z\}/\operatorname{Jac} F < \infty$.
- Embedded topology: topological type of $(\mathbb{S}^5, K)$.
- Main tool, Milnor fibration $(0 < \eta \ll \varepsilon < \varepsilon_0)$:

$$\mathbb{S}^5 \setminus K \xrightarrow{\frac{f}{|f|}} \mathbb{S}^1 \qquad\qquad \overline{\mathbb{B}}^6_\varepsilon \cap f^{-1}(\mathbb{S}^1_\eta) \xrightarrow{f} \mathbb{S}^1_\eta$$

- Milnor fiber $F \simeq \bigvee_\mu \mathbb{S}^2$.
- Fibrations determined by $\rho : F \to F$, diffeomorphism well-defined up to isotopy (monodromy).
- Homological monodromy: $\rho^* : H^2(F; R) \to H^2(F; R)$, $R = \mathbb{Z}, \mathbb{Q}, \mathbb{C}$.
- Intersection and Seifert forms on $H_2(F; \mathbb{Z})$ (distinguished basis of vanishing cycles)



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.

Total and strict transform, A'Campo's formula

- $\hat{S} := \overline{\pi^{-1}(S) \setminus \{0\}}$, strict transform, $\pi_{|\hat{S}}$ is a good resolution.



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.

Total and strict transform, A'Campo's formula

- $\hat{S} := \overline{\pi^{-1}(S) \setminus \{0\}}$, strict transform, $\pi_{|\hat{S}}$ is a good resolution.
- $\pi^* S = \sum_{i \in I} m_i E_i$, $0 \in I$, $(E_0, m_0) = (\hat{S}, 1)$, total transform.



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.

Total and strict transform, A'Campo's formula

- $\hat{S} := \overline{\pi^{-1}(S) \setminus \{0\}}$, strict transform, $\pi_{|\hat{S}}$ is a good resolution.
- $\pi^* S = \sum_{i \in I} m_i E_i$, $0 \in I$, $(E_0, m_0) = (\hat{S}, 1)$, total transform.
- $\check{E}_J := \bigcap_{i \in J} E_i \setminus \bigcup_{i \in I \setminus J} E_i$, $J \subset I$.



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.

Total and strict transform, A'Campo's formula

- $\hat{S} := \overline{\pi^{-1}(S) \setminus \{0\}}$, strict transform, $\pi_{|\hat{S}}$ is a good resolution.
- $\pi^* S = \sum_{i \in I} m_i E_i$, $0 \in I$, $(E_0, m_0) = (\hat{S}, 1)$, total transform.
- $\check{E}_J := \bigcap_{i \in J} E_i \setminus \bigcup_{i \in I \setminus J} E_i$, $J \subset I$.
- $\zeta(t) = \prod_{k=0}^{2} \Delta^{\rho}_{H^k(F;\mathbb{C})}(t)^{(-1)^k} = (t-1)\Delta^{\rho}_{H^2(F;\mathbb{C})}(t) = \prod_{i \in I} (t^{m_i} - 1)^{\chi(\check{E}_i)}$.



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.

Example

- $F = x^d + y^d + z^d$, cone over a smooth curve C_d
- $\hat{S}$ cylinder over C_d



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.

Example

- $F = x^d + y^d + z^d$, cone over a smooth curve C_d
- $\hat{S}$ cylinder over C_d
- $I = \{0,1\}$, $E_1 \cong \mathbb{P}^2$, $m_1 = d$, $E_0 \cap E_1 = C_d$



Embedded resolution

Definition

An *embedded resolution* of $(S,0) \subset (\mathbb{C}^3,0)$ is a proper birational map $\pi : (X,E) \to (\mathbb{C}^3,0)$, $E = \pi^{-1}(0)$, such that π is an isomorphism on $X \setminus E$ and $\pi^{-1}(S)$ is a normal crossing divisor.

Example

- $F = x^d + y^d + z^d$, cone over a smooth curve C_d
- $\hat{S}$ cylinder over C_d
- $I = \{0,1\}$, $E_1 \cong \mathbb{P}^2$, $m_1 = d$, $E_0 \cap E_1 = C_d$
- $\Delta^{\rho}_{H^2(F;\mathbb{C})}(t) = \dfrac{(t^d-1)^{d^2-3d+3}}{t-1}$.
- Since it is homogeneous, the monodromy is semisimple (the geometric monodromy has finite order d).



Semistable resolution and Steenbrink spectral sequence

- $e = \operatorname{lcm}(m_i \mid i \in I)$, ν normalization, $D_0 \equiv \hat{S}$, D_i ramified cover over E_i.

$$\begin{array}{ccccc} (Z, D) & \xrightarrow{\nu} & (\tilde{X}, E) & \xrightarrow{\tau_e} & (X, E) \\ & & \downarrow & & \downarrow F \circ \pi \\ \text{Model for the} & & (\mathbb{C}, 0) & \longrightarrow & (\mathbb{C}, 0) \\ \text{Milnor fibration} & & t & \longmapsto & t^e \end{array}$$



Semistable resolution and Steenbrink spectral sequence

- Steenbrink spectral sequence degenerates at E_2, converges to $H^*(F;\mathbb{C})$ filtered by the weight filtration. E_1 step:

$$D^{[0]}:=\coprod_{i\in I_0} D_i, \quad D^{[1]}:=\coprod_{I_0} D_i\cap D_j, \quad D_0^{[1]}:=\coprod_{I} D_i\cap D_j$$
$$D^{[2]}:=\coprod_{I_0} D_i\cap D_j\cap D_k, \quad D_0^{[2]}:=\coprod_{I} D_i\cap D_j\cap D_k, \quad I_0:=I\setminus\{0\}$$

$$\begin{array}{ccccccc}
H^0(D_0^{[2]};\mathbb{C}) & & & & & & \\
\uparrow{\scriptstyle i^*} & & & & & & \\
H^0(D^{[1]};\mathbb{C}) & H^1(D^{[1]};\mathbb{C}) & & H^2(D^{[1]};\mathbb{C}) & & & \\
\uparrow{\scriptstyle i^*} & \uparrow{\scriptstyle i^*} & & \uparrow{\scriptstyle i^*\oplus g} & & & \\
H^0(D^{[0]};\mathbb{C}) & H^1(D^{[0]};\mathbb{C}) & H^2(D^{[0]};\mathbb{C})\oplus H^0(D_0^{[2]};\mathbb{C}) & H^3(D^{[0]};\mathbb{C}) & H^4(D^{[0]};\mathbb{C}) \\
 & & \uparrow{\scriptstyle g} & \uparrow{\scriptstyle g} & \uparrow{\scriptstyle g} \\
 & & H^0(D_0^{[1]};\mathbb{C}) & H^1(D_0^{[1]};\mathbb{C}) & H^2(D_0^{[1]};\mathbb{C}) \\
 & & & & \uparrow{\scriptstyle g} \\
 & & & & H^0(D_0^{[2]};\mathbb{C})
\end{array}$$



Semistable resolution and Steenbrink spectral sequence

- Δ_D, Δ_E, dual complexes of the normal crossing divisors D, E, $\pi_\Delta : \Delta_D \to \Delta_E$, $\rho_\Delta : \Delta_D \to \Delta_D$ the monodromy.
  - Δ_D is connected, $H^1(\Delta_D; \mathbb{C}) = 0$.
  - The characteristic polynomial of ρ_Δ on $H^2(\Delta_D)$ measures the Jordan blocks of size 3.
  - It corresponds to the first column.
- Second column: the characteristic polynomial of the monodromy acting on $H^1(D^{[1]}; \mathbb{C})/H^1(D^{[0]}; \mathbb{C})$ measures the Jordan blocks of size 2 (and eigenvalue $\neq 1$).



Semistable resolution and Steenbrink spectral sequence

- Δ_D, Δ_E, dual complexes of the normal crossing divisors D, E, $\pi_\Delta : \Delta_D \to \Delta_E$, $\rho_\Delta : \Delta_D \to \Delta_D$ the monodromy.
  - Δ_D is connected, $H^1(\Delta_D; \mathbb{C}) = 0$.
  - The characteristic polynomial of ρ_Δ on $H^2(\Delta_D)$ measures the Jordan blocks of size 3.
  - It corresponds to the first column.
- Second column: the characteristic polynomial of the monodromy acting on $H^1(D^{[1]}; \mathbb{C})/H^1(D^{[0]}; \mathbb{C})$ measures the Jordan blocks of size 2 (and eigenvalue $\neq 1$).



Semistable resolution and Steenbrink spectral sequence

- Δ_D, Δ_E, dual complexes of the normal crossing divisors D, E, $\pi_\Delta : \Delta_D \to \Delta_E$, $\rho_\Delta : \Delta_D \to \Delta_D$ the monodromy.
  - Δ_D is connected, $H^1(\Delta_D; \mathbb{C}) = 0$.
  - The characteristic polynomial of ρ_Δ on $H^2(\Delta_D)$ measures the Jordan blocks of size 3.
    - It corresponds to the first column.
- Second column: the characteristic polynomial of the monodromy acting on $H^1(D^{[1]}; \mathbb{C})/H^1(D^{[0]}; \mathbb{C})$ measures the Jordan blocks of size 2 (and eigenvalue $\neq 1$).



Semistable resolution and Steenbrink spectral sequence

- Δ_D, Δ_E, dual complexes of the normal crossing divisors D, E, $\pi_\Delta : \Delta_D \to \Delta_E$, $\rho_\Delta : \Delta_D \to \Delta_D$ the monodromy.
  - Δ_D is connected, $H^1(\Delta_D; \mathbb{C}) = 0$.
  - The characteristic polynomial of ρ_Δ on $H^2(\Delta_D)$ measures the Jordan blocks of size 3.
  - It corresponds to the first column.
- Second column: the characteristic polynomial of the monodromy acting on $H^1(D^{[1]}; \mathbb{C})/H^1(D^{[0]}; \mathbb{C})$ measures the Jordan blocks of size 2 (and eigenvalue $\neq 1$).



Semistable resolution and Steenbrink spectral sequence

- Δ_D, Δ_E, dual complexes of the normal crossing divisors D, E, $\pi_\Delta : \Delta_D \to \Delta_E$, $\rho_\Delta : \Delta_D \to \Delta_D$ the monodromy.
  - Δ_D is connected, $H^1(\Delta_D; \mathbb{C}) = 0$.
  - The characteristic polynomial of ρ_Δ on $H^2(\Delta_D)$ measures the Jordan blocks of size 3.
  - It corresponds to the first column.
- Second column: the characteristic polynomial of the monodromy acting on $H^1(D^{[1]}; \mathbb{C})/H^1(D^{[0]}; \mathbb{C})$ measures the Jordan blocks of size 2 (and eigenvalue $\neq 1$).



Surfaces in $\mathbb{C}^3$

Superisolated and Lê-Yomdin singularities

Zariski tuples and Jordan form

Zariski tuples of non-Alexander type

Quasi-homogeneous and Weighted Lê-Yomdin singularities



Curves and homogeneous singularities

Homogeneous polynomials

$F(x, y, z) \in \mathbb{C}[x, y, z]$ homogeneous polynomial of degree $d > 0$:

- Projective curve in $\mathbb{P}^2$.
- Non-isolated (in general) singularity in $\mathbb{C}^3$.



Curves and homogeneous singularities

Homogeneous polynomials

$F(x,y,z) \in \mathbb{C}[x,y,z]$ homogeneous polynomial of degree $d > 0$:

- Projective curve in $\mathbb{P}^2$.
- Non-isolated (in general) singularity in $\mathbb{C}^3$.

Superisolated singularities: definition 1

$(S,0) = \{f(x,y,z) = 0\}$, is *superisolated* (SIS) if $\mathrm{Sing}(C_m) \cap C_{m+1} = \varnothing$,

$$f(x,y,z) = \overbrace{f_m(x,y,z)}^{\neq 0} + f_{m+1}(x,y,z) + \cdots \in \mathbb{C}\{x,y,z\},$$

$C_j = V_{\mathbb{P}}(f_j) \subset \mathbb{P}^2$, C_m the tangent cone.



Curves and homogeneous singularities

Homogeneous polynomials

$F(x,y,z) \in \mathbb{C}[x,y,z]$ homogeneous polynomial of degree $d > 0$:

- Projective curve in $\mathbb{P}^2$.
- Non-isolated (in general) singularity in $\mathbb{C}^3$.

Superisolated singularities: definition 1

$(S,0) = \{f(x,y,z) = 0\}$, is *superisolated* (SIS) if $\mathrm{Sing}(C_m) \cap C_{m+1} = \varnothing$,

$$f(x,y,z) = \overbrace{f_m(x,y,z)}^{\neq 0} + f_{m+1}(x,y,z) + \cdots \in \mathbb{C}\{x,y,z\},$$

$C_j = V_{\mathbb{P}}(f_j) \subset \mathbb{P}^2$, C_m the tangent cone.

- Generic deformation of $f_m(x,y,z) = 0$ to be isolated singularity.
- C_m must be reduced.



Curves and homogeneous singularities

Homogeneous polynomials

$F(x, y, z) \in \mathbb{C}[x, y, z]$ homogeneous polynomial of degree $d > 0$:

- Projective curve in $\mathbb{P}^2$.
- Non-isolated (in general) singularity in $\mathbb{C}^3$.

Superisolated singularities: definition 2

The germ $(S, 0)$ is *superisolated* (SIS) if the strict transform of S under the blow-up of the origin is smooth.



Curves and homogeneous singularities

Homogeneous polynomials

$F(x,y,z) \in \mathbb{C}[x,y,z]$ homogeneous polynomial of degree $d > 0$:

- Projective curve in $\mathbb{P}^2$.
- Non-isolated (in general) singularity in $\mathbb{C}^3$.

Superisolated singularities: definition 2

The germ $(S,0)$ is *superisolated* (SIS) if the strict transform of S under the blow-up of the origin is smooth.

Lê-Yomdin singularities

The germ $(S,0)$ is *Lê-Yomdin* if $f_{m+1} = \cdots = f_{m+k-1} = 0$ and $\mathrm{Sing}(C_m) \cap C_{m+k} = \varnothing$.



Curves and homogeneous singularities

Homogeneous polynomials

$F(x, y, z) \in \mathbb{C}[x, y, z]$ homogeneous polynomial of degree $d > 0$:

- Projective curve in $\mathbb{P}^2$.
- Non-isolated (in general) singularity in $\mathbb{C}^3$.

Superisolated singularities: definition 2

The germ $(S, 0)$ is *superisolated* (SIS) if the strict transform of S under the blow-up of the origin is smooth.

Lê-Yomdin singularities

The germ $(S, 0)$ is *Lê-Yomdin* if $f_{m+1} = \cdots = f_{m+k-1} = 0$ and $\mathrm{Sing}(C_m) \cap C_{m+k} = \varnothing$.
If the tangent cone is not smooth, the strict transform under the blow-up has k-cyclic singularities.




Examples and results

- $S = \{F = 0\}$, $F = y^2z - x^3 - z^4$.



Examples and results


- $S = \{F = 0\}$, $F = y^2z - x^3 - z^4$.
- Blow-up the origin of $\mathbb{C}^3$:





Examples and results


- $S = \{F = 0\}$, $F = y^2z - x^3 - z^4$.
- Blow-up the origin of $\mathbb{C}^3$:






Examples and results

- $S = \{F = 0\}$, $F = y^2z - x^3 - z^4$.
- Blow-up the origin of $\mathbb{C}^3$:




Examples and results

- $S = \{F = 0\}$, $F = y^2z - x^3 - z^4$.
- Abstract resolution and good resolution:



Examples and results

- $S = \{F = 0\}$, $F = y^2z - x^3 - z^4$.
- Abstract resolution and good resolution:



Theorem (I. Luengo)


S SIS with tangent cone $C_1 + \cdots + C_r$, $\deg C_i = d_i$. Then C in $\hat{S}$ is abstractly isomorphic to C in $\mathbb{P}^2$ and

$$(C_i \cdot C_i)_{\hat{S}} = -d_i(d - d_i + 1), \quad (C_i \cdot C_j)_{\hat{S}} = d_i d_j (i \neq j).$$



Examples and results

- $S = \{F = 0\}$, $F = y^2z - x^3 - z^4$.
- Abstract resolution and good resolution:



Theorem (I. Luengo)

S SIS with tangent cone $C_1 + \cdots + C_r$, $\deg C_i = d_i$. Then C in $\hat{S}$ is abstractly isomorphic to C in $\mathbb{P}^2$ and

$$(C_i \cdot C_i)_{\hat{S}} = -d_i(d - d_i + 1), \quad (C_i \cdot C_j)_{\hat{S}} = d_i d_j (i \neq j).$$

Theorem (J. Stevens, E. A.)

S SIS with tangent cone C of degree d:

$$\Delta^{\rho}_{H^2(F;\mathbb{C})} = \frac{(t^d - 1)^{d^2 - 3d + 3 - \mu(C)}}{t - 1} \prod_{P \in \mathrm{Sing}(C)} \Delta^{\rho_P}_{H^1(F_p;\mathbb{C})}(t^{d+1}).$$



Yomdin singularities

Theorem (E. A., J.I. Cogolludo, J. Martín-Morales)

S k-Lê-Yomdin singularity with tangent cone $C_1 + \cdots + C_r$, $\deg C_i = d_i$. Then C in $\hat{S}$ is abstractly isomorphic to C in $\mathbb{P}^2$, has cyclic singularities at $\mathrm{Sing}(C)$ *and*

$$(C_i \cdot C_i)_{\hat{S}} = -\frac{d_i(d - d_i + k)}{k}, \quad (C_i \cdot C_j)_{\hat{S}} = \frac{d_i d_j}{k} (i \neq j).$$



Yomdin singularities

Theorem (E. A., J.I. Cogolludo, J. Martín-Morales)

S k-Lê-Yomdin singularity with tangent cone $C_1 + \cdots + C_r$, $\deg C_i = d_i$. Then C in $\hat{S}$ is abstractly isomorphic to C in $\mathbb{P}^2$, has cyclic singularities at $\mathrm{Sing}(C)$ *and*

$$(C_i \cdot C_i)_{\hat{S}} = -\frac{d_i(d - d_i + k)}{k}, \quad (C_i \cdot C_j)_{\hat{S}} = \frac{d_i d_j}{k} (i \neq j).$$

Theorem (S. Gussein-Zade-I. Luengo-A. Melle, J. Martín)

S k-Lê-Yomdin singularity with tangent cone C of degree d:

$$\Delta^{\rho}_{H^2(F;\mathbb{C})} = \frac{(t^d - 1)^{d^2 - 3d + 3 - \mu(C)}}{t - 1} \prod_{P \in \mathrm{Sing}(C)} \Delta^{\rho_P,(k)}_{H^1(F_p;\mathbb{C})}(t^{d+k}).$$

where

$$\Delta(t) = \prod_{j=1}^{r} (t^{m_j} - 1)^{n_j} \Longrightarrow \Delta^{(k)}(t) = \prod_{j=1}^{r} \left(t^{\frac{m_j}{\gcd(m_j,k)}} - 1 \right)^{\gcd(m_j,k) n_j}$$



Surfaces in $\mathbb{C}^3$

Superisolated and Lê-Yomdin singularities

Zariski tuples and Jordan form

Zariski tuples of non-Alexander type

Quasi-homogeneous and Weighted Lê-Yomdin singularities



Combinatorics

Yau's conjecture

$(S,0) \subset (\mathbb{C}^3,0)$ isolated surface singularity. Then the embedded topological type of $(S,0)$ determines and is determined by $\Delta^{\rho}_{H^2(F;\mathbb{C})}$ and $\pi_1(K)$ (equivalent to the topological type of K).



Combinatorics

Yau's conjecture

$(S,0) \subset (\mathbb{C}^3,0)$ isolated surface singularity. Then the embedded topological type of $(S,0)$ determines and is determined by $\Delta^{\rho}_{H^2(F;\mathbb{C})}$ and $\pi_1(K)$ (equivalent to the topological type of K).

Definition

$C \subset \mathbb{P}^2$ reduced. The *combinatorics* of C is composed by the following data:



Combinatorics

Yau's conjecture

$(S,0) \subset (\mathbb{C}^3,0)$ isolated surface singularity. Then the embedded topological type of $(S,0)$ determines and is determined by $\Delta^{\rho}_{H^2(F;\mathbb{C})}$ and $\pi_1(K)$ (equivalent to the topological type of K).

Definition

$C \subset \mathbb{P}^2$ reduced. The *combinatorics* of C is composed by the following data:

- degrees of irreducible components,



Combinatorics

Yau's conjecture

$(S,0) \subset (\mathbb{C}^3,0)$ isolated surface singularity. Then the embedded topological type of $(S,0)$ determines and is determined by $\Delta^{\rho}_{H^2(F;\mathbb{C})}$ and $\pi_1(K)$ (equivalent to the topological type of K).

Definition

$C \subset \mathbb{P}^2$ reduced. The *combinatorics* of C is composed by the following data:

- degrees of irreducible components,
- topological type of singularities, and



Combinatorics

Yau's conjecture

$(S,0) \subset (\mathbb{C}^3,0)$ isolated surface singularity. Then the embedded topological type of $(S,0)$ determines and is determined by $\Delta^{\rho}_{H^2(F;\mathbb{C})}$ and $\pi_1(K)$ (equivalent to the topological type of K).

Definition

$C \subset \mathbb{P}^2$ reduced. The *combinatorics* of C is composed by the following data:

- degrees of irreducible components,
- topological type of singularities, and
- assignment of branches of the singular points to irreducible components.



Combinatorics

Yau's conjecture

$(S,0) \subset (\mathbb{C}^3,0)$ isolated surface singularity. Then the embedded topological type of $(S,0)$ determines and is determined by $\Delta^{\rho}_{H^2(F;\mathbb{C})}$ and $\pi_1(K)$ (equivalent to the topological type of K).

Definition

$C \subset \mathbb{P}^2$ reduced. The *combinatorics* of C is composed by the following data:

- degrees of irreducible components,
- topological type of singularities, and
- assignment of branches of the singular points to irreducible components.

Corollary

$(S_i,0)$, $i=1,2$, SISs with tangent cones having the same combinatorics. They have the same characteristic polynomial of the monodromy and the same embedded topology.



Zariski tuples

Definition

$C_1, \ldots, C_r \subset \mathbb{P}^2$ curves with the same combinatorics. They form a *Zariski r-tuple* if $(\mathbb{P}^2, C_i)$ and $(\mathbb{P}^2, C_j)$ are not homeomorphic if $i \neq j$.



Zariski tuples

Definition

$C_1, \ldots, C_r \subset \mathbb{P}^2$ curves with the same combinatorics. They form a *Zariski r-tuple* if $(\mathbb{P}^2, C_i)$ and $(\mathbb{P}^2, C_j)$ are not homeomorphic if $i \neq j$.

Problem

Let $(S_i, 0)$, $i = 1, \ldots, r$, SISs such that their tangent cones form a Zariski tuple. How can we compare the embedded topology of the SISs?



Zariski tuples

Definition

$C_1, \ldots, C_r \subset \mathbb{P}^2$ curves with the same combinatorics. They form a *Zariski r-tuple* if $(\mathbb{P}^2, C_i)$ and $(\mathbb{P}^2, C_j)$ are not homeomorphic if $i \neq j$.

Problem

Let $(S_i, 0)$, $i = 1, \ldots, r$, SISs such that their tangent cones form a Zariski tuple. How can we compare the embedded topology of the SISs?

Definition (Libgober)

$C \subset \mathbb{P}^2$ reduced curve of degree d. The *Alexander polynomial* Δ_C of C is the characteristic polynomial of the monodromy on H^1 of the branched cover of $\mathbb{P}^2$ ramified along C.



Zariski tuples

Definition

$C_1, \ldots, C_r \subset \mathbb{P}^2$ curves with the same combinatorics. They form a *Zariski r-tuple* if $(\mathbb{P}^2, C_i)$ and $(\mathbb{P}^2, C_j)$ are not homeomorphic if $i \neq j$.

Problem

Let $(S_i, 0)$, $i = 1, \ldots, r$, SISs such that their tangent cones form a Zariski tuple. How can we compare the embedded topology of the SISs?

Definition (Libgober)

$C \subset \mathbb{P}^2$ reduced curve of degree d. The *Alexander polynomial* Δ_C of C is the characteristic polynomial of the monodromy on H^1 of the branched cover of $\mathbb{P}^2$ ramified along C.

Definition

A Zariski r-tuple is an *Alexander-Zariski r-tuple* if the Alexander polynomials are pairwise distinct.



Alexander-Zariski tuples polynomials and SIS

Alexander-Zariski polynomials exist (Zariski)

C sextic, $\#\,\mathrm{Sing}(C) = 6$, ordinary cusps.

$$\Delta(t) = \begin{cases} t^2 - t + 1 & \text{if } \mathrm{Sing}\, C \subset \text{ a conic} \\ 1 & \text{otherwise} \end{cases}$$



Alexander-Zariski tuples polynomials and SIS

Alexander-Zariski polynomials exist (Zariski)

C sextic, $\#\operatorname{Sing}(C) = 6$, ordinary cusps.

$$\Delta(t) = \begin{cases} t^2 - t + 1 & \text{if } \operatorname{Sing} C \subset \text{ a conic} \\ 1 & \text{otherwise} \end{cases}$$

2-Jordan blocks for SIS

$D^{[0]} = D_o \coprod D_o^{[0]}$, where D_o is birationally equivalent to the d-cyclic cover of $\mathbb{P}^2$ ramified along the tangent cone. The number of 2-Jordan blocks (eigenvalues $\neq 1$):

$$\underbrace{\dim_{\mathbb{C}} H^1(D^{[1]};\mathbb{C}) - \dim_{\mathbb{C}} H^1(D_0^{[0]};\mathbb{C})}_{\text{combinatorial}} - \underbrace{\dim_{\mathbb{C}} H^1(D_o;\mathbb{C})}_{\deg \Delta_C}$$



Alexander-Zariski tuples polynomials and SIS

Alexander-Zariski polynomials exist (Zariski)

C sextic, $\#\operatorname{Sing}(C) = 6$, ordinary cusps.

$$\Delta(t) = \begin{cases} t^2 - t + 1 & \text{if } \operatorname{Sing} C \subset \text{ a conic} \\ 1 & \text{otherwise} \end{cases}$$

2-Jordan blocks for SIS

$D^{[0]} = D_o \coprod D_o^{[0]}$, where D_o is birationally equivalent to the d-cyclic cover of $\mathbb{P}^2$ ramified along the tangent cone. The number of 2-Jordan blocks (eigenvalues $\neq 1$):

$$\underbrace{\dim_{\mathbb{C}} H^1(D^{[1]};\mathbb{C}) - \dim_{\mathbb{C}} H^1(D_0^{[0]};\mathbb{C})}_{\text{combinatorial}} - \underbrace{\dim_{\mathbb{C}} H^1(D_o;\mathbb{C})}_{\deg \Delta_C}$$

Corollary

Zariski's example provides a counterexample to Yau's conjecture.



Surfaces in $\mathbb{C}^3$

Superisolated and Lê-Yomdin singularities

Zariski tuples and Jordan form


Zariski tuples of non-Alexander type

Quasi-homogeneous and Weighted Lê-Yomdin singularities



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,
- $\mathbf{n} := (n_1, \ldots, n_r) \in (\mathbb{Z}_{>0})^r$, coprime



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,
- $\mathbf{n} := (n_1, \ldots, n_r) \in (\mathbb{Z}_{>0})^r$, coprime
- $C_{\mathbf{n}} := n_1 C_1 + \cdots + n_r C_r$, $d = n_1 + \cdots + n_r$.



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,
- $\mathbf{n} := (n_1, \ldots, n_r) \in (\mathbb{Z}_{>0})^r$, coprime
- $C_{\mathbf{n}} := n_1 C_1 + \cdots + n_r C_r$, $d = n_1 + \cdots + n_r$.
- **n**-Alexander polynomial associated to the unramified cyclic d-cover of $\mathbb{P}^2 \setminus C_{\mathbf{n}}$.



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,
- $\mathbf{n} := (n_1, \ldots, n_r) \in (\mathbb{Z}_{>0})^r$, coprime
- $C_{\mathbf{n}} := n_1 C_1 + \cdots + n_r C_r$, $d = n_1 + \cdots + n_r$.
- $\mathbf{n}$-Alexander polynomial associated to the unramified cyclic d-cover of $\mathbb{P}^2 \setminus C_{\mathbf{n}}$.
- Characteristic varieties measure these Alexander polynomials all together



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,
- $\mathbf{n} := (n_1, \ldots, n_r) \in (\mathbb{Z}_{>0})^r$, coprime
- $C_{\mathbf{n}} := n_1 C_1 + \cdots + n_r C_r$, $d = n_1 + \cdots + n_r$.
- $\mathbf{n}$-Alexander polynomial associated to the unramified cyclic d-cover of $\mathbb{P}^2 \setminus C_{\mathbf{n}}$.
- Characteristic varieties measure these Alexander polynomials all together
- There are Zariski tuples not of Alexander type, distinguished by characteristic varieites.



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,
- $\mathbf{n} := (n_1, \ldots, n_r) \in (\mathbb{Z}_{>0})^r$, coprime
- $C_{\mathbf{n}} := n_1 C_1 + \cdots + n_r C_r$, $d = n_1 + \cdots + n_r$.
- $\mathbf{n}$-Alexander polynomial associated to the unramified cyclic d-cover of $\mathbb{P}^2 \setminus C_{\mathbf{n}}$.
- Characteristic varieties measure these Alexander polynomials all together
- There are Zariski tuples not of Alexander type, distinguished by characteristic varieites.
- What about the embedded topology of the associated SIS or k-Lê-Yomdin?



Characteristic varieties

- $C = C_1 + \cdots + C_r$, $\deg C_i = d_i$,
- $\mathbf{n} := (n_1, \ldots, n_r) \in (\mathbb{Z}_{>0})^r$, coprime
- $C_{\mathbf{n}} := n_1 C_1 + \cdots + n_r C_r$, $d = n_1 + \cdots + n_r$.
- $\mathbf{n}$-Alexander polynomial associated to the unramified cyclic d-cover of $\mathbb{P}^2 \setminus C_{\mathbf{n}}$.
- Characteristic varieties measure these Alexander polynomials all together
- There are Zariski tuples not of Alexander type, distinguished by characteristic varieites.
- What about the embedded topology of the associated SIS or k-Lê-Yomdin?
- One may construct *generic isolated singularities* with non-reduced tangent cone with distinct embedded topology.



Fundamental group

- There are Zariski pairs whose only tractable invariant to distinguish them is the fundamental group of the complement (e.g, Rybnikov line arrangements).



Fundamental group

- There are Zariski pairs whose only tractable invariant to distinguish them is the fundamental group of the complement (e.g, Rybnikov line arrangements).
- From semistable normalization, the unramified cyclic cover of $\mathbb{P}^2 \setminus C$ is the main part of a decomposition of the Milnor fiber F (kind of Thurston-Nielsen decomposition), but F is simply connected!



Fundamental group

- There are Zariski pairs whose only tractable invariant to distinguish them is the fundamental group of the complement (e.g, Rybnikov line arrangements).
- From semistable normalization, the unramified cyclic cover of $\mathbb{P}^2 \setminus C$ is the main part of a decomposition of the Milnor fiber F (kind of Thurston-Nielsen decomposition), but F is simply connected!
- There are also Zariski tuples with isomorphic fundamental group, even abelian fundamental group.



Fundamental group

- There are Zariski pairs whose only tractable invariant to distinguish them is the fundamental group of the complement (e.g, Rybnikov line arrangements).
- From semistable normalization, the unramified cyclic cover of $\mathbb{P}^2 \setminus C$ is the main part of a decomposition of the Milnor fiber F (kind of Thurston-Nielsen decomposition), but F is simply connected!
- There are also Zariski tuples with isomorphic fundamental group, even abelian fundamental group.
- Finally, there are *arithmetic Zariski tuples*, i.e., a curve defined over a number field $\mathbb{K}$ and the elements of the tuple correspond to distinct embeddings $\mathbb{K} \hookrightarrow \mathbb{C}$.



Fundamental group

- There are Zariski pairs whose only tractable invariant to distinguish them is the fundamental group of the complement (e.g, Rybnikov line arrangements).
- From semistable normalization, the unramified cyclic cover of $\mathbb{P}^2 \setminus C$ is the main part of a decomposition of the Milnor fiber F (kind of Thurston-Nielsen decomposition), but F is simply connected!
- There are also Zariski tuples with isomorphic fundamental group, even abelian fundamental group.
- Finally, there are *arithmetic Zariski tuples*, i.e., a curve defined over a number field $\mathbb{K}$ and the elements of the tuple correspond to distinct embeddings $\mathbb{K} \hookrightarrow \mathbb{C}$.
- Same question: What about the embedded topology of the associated SIS or k-Lê-Yomdin?



Intersection form (Arnol'd-Varchenko-Gussein-Zade)

- $\cdot_\partial : H_2(F, \partial F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, duality pairing,

  $D : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})^*, \quad D^* : H_2(F; \mathbb{Z}) \to H_2(F, \partial F; \mathbb{Z})^*.$



Intersection form (Arnol'd-Varchenko-Gussein-Zade)

- $\cdot_\partial : H_2(F, \partial F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, duality pairing,

  $D : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})^*, \quad D^* : H_2(F; \mathbb{Z}) \to H_2(F, \partial F; \mathbb{Z})^*.$

- $\exists \rho : F \to F$ monodromy, $\rho_{|\partial F} = \mathbf{1}_{\partial F}$,
  $\mathrm{var} : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})$, on cycles $\rho - \mathbf{1}$, *variation isomorphism.*



Intersection form (Arnol'd-Varchenko-Gussein-Zade)

- $\cdot_\partial : H_2(F, \partial F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, duality pairing,

  $D : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})^*, \quad D^* : H_2(F; \mathbb{Z}) \to H_2(F, \partial F; \mathbb{Z})^*.$

- $\exists \rho : F \to F$ monodromy, $\rho_{|\partial F} = \mathbf{1}_{\partial F}$,
  $\mathrm{var} : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})$, on cycles $\rho - \mathbf{1}$, *variation isomorphism.*
- $L : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $L(a, b) := \mathrm{var}^{-1}(a) \cdot_\partial b$.



Intersection form (Arnol'd-Varchenko-Gussein-Zade)

- $\cdot_\partial : H_2(F, \partial F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, duality pairing,

  $$D : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})^*, \quad D^* : H_2(F; \mathbb{Z}) \to H_2(F, \partial F; \mathbb{Z})^*.$$

- $\exists \rho : F \to F$ monodromy, $\rho_{|\partial F} = \mathbf{1}_{\partial F}$,
  $\mathrm{var} : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})$, on cycles $\rho - \mathbf{1}$, *variation isomorphism.*
- $L : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $L(a, b) := \mathrm{var}^{-1}(a) \cdot_\partial b$.
- L is the Seifert form.



Intersection form (Arnol'd-Varchenko-Gussein-Zade)

- $\cdot_\partial : H_2(F, \partial F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, duality pairing,

  $$D : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})^*, \quad D^* : H_2(F; \mathbb{Z}) \to H_2(F, \partial F; \mathbb{Z})^*.$$

- $\exists \rho : F \to F$ monodromy, $\rho_{|\partial F} = \mathbf{1}_{\partial F}$,
  $\mathrm{var} : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})$, on cycles $\rho - \mathbf{1}$, *variation isomorphism.*
- $L : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $L(a, b) := \mathrm{var}^{-1}(a) \cdot_\partial b$.
- L is the Seifert form.
- $\cdot : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $a \cdot b = -L(a, b) - L(b, a)$.



Intersection form (Arnol'd-Varchenko-Gussein-Zade)

- $\cdot_\partial : H_2(F, \partial F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, duality pairing,

  $$D : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})^*, \quad D^* : H_2(F; \mathbb{Z}) \to H_2(F, \partial F; \mathbb{Z})^*.$$

- $\exists \rho : F \to F$ monodromy, $\rho_{|\partial F} = \mathbf{1}_{\partial F}$,
  $\mathrm{var} : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})$, on cycles $\rho - \mathbf{1}$, *variation isomorphism.*
- $L : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $L(a, b) := \mathrm{var}^{-1}(a) \cdot_\partial b$.
- L is the Seifert form.
- $\cdot : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $a \cdot b = -L(a, b) - L(b, a)$.
- Transposed variation

$$\begin{array}{ccc} H_2(F, \partial F; \mathbb{Z}) & \xrightarrow{{}^t\mathrm{var}} & H_2(F; \mathbb{Z}) \\ \downarrow D & & \downarrow D^* \\ H_2(F; \mathbb{Z})^* & \xrightarrow{\mathrm{var}^*} & H_2(F, \partial F; \mathbb{Z})^* \end{array}$$



Intersection form (Arnol'd-Varchenko-Gussein-Zade)

- $\cdot_\partial : H_2(F, \partial F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, duality pairing,

  $$D : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})^*, \quad D^* : H_2(F; \mathbb{Z}) \to H_2(F, \partial F; \mathbb{Z})^*.$$

- $\exists \rho : F \to F$ monodromy, $\rho_{|\partial F} = \mathbf{1}_{\partial F}$,
  $\mathrm{var} : H_2(F, \partial F; \mathbb{Z}) \to H_2(F; \mathbb{Z})$, on cycles $\rho - \mathbf{1}$, *variation isomorphism.*
- $L : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $L(a, b) := \mathrm{var}^{-1}(a) \cdot_\partial b$.
- L is the Seifert form.
- $\cdot : H_2(F; \mathbb{Z}) \times H_2(F; \mathbb{Z}) \to \mathbb{Z}$, $a \cdot b = -L(a, b) - L(b, a)$.
- Transposed variation

$$\begin{array}{ccc} H_2(F, \partial F; \mathbb{Z}) & \xrightarrow{{}^t\mathrm{var}} & H_2(F; \mathbb{Z}) \\ \downarrow D & & \downarrow D^* \\ H_2(F; \mathbb{Z})^* & \xrightarrow{\mathrm{var}^*} & H_2(F, \partial F; \mathbb{Z})^* \end{array}$$

- $\rho_* : H_2(F; \mathbb{Z}) \to H_2(F; \mathbb{Z})$, $\rho_* = -\mathrm{var} \circ {}^t\mathrm{var}^{-1}$.



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

$(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.
- Var upper triangular with diagonals 1, Vart = ^{t}Var.



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.
- Var upper triangular with diagonals 1, Vart = ^{t}Var.
- $S = {}^t\mathrm{Var}^{-1}$, $I = -{}^t\mathrm{Var}^{-1} - \mathrm{Var}^{-1}$, $M = -\,\mathrm{Var}\,{}^t\mathrm{Var}^{-1}$.



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.
- Var upper triangular with diagonals 1, Vart = ^{t}Var.
- $S = {}^t\mathrm{Var}^{-1}$, $I = -{}^t\mathrm{Var}^{-1} - \mathrm{Var}^{-1}$, $M = -\,\mathrm{Var}\,{}^t\mathrm{Var}^{-1}$.
- Gabriélov polar method and Escario's give I for a DVBC. A'Campo can work with integral bilinear forms of high rank.



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.
- Var upper triangular with diagonals 1, Vart = ^{t}Var.
- $S = {}^t\mathrm{Var}^{-1}$, $I = -{}^t\mathrm{Var}^{-1} - \mathrm{Var}^{-1}$, $M = -\,\mathrm{Var}\,{}^t\mathrm{Var}^{-1}$.
- Gabriélov polar method and Escario's give I for a DVBC. A'Campo can work with integral bilinear forms of high rank.

A'Campo's question

Is ρ_* conjugated to its inverse (over $\mathbb{Z}$)?



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.
- Var upper triangular with diagonals 1, Vart = ^{t}Var.
- $S = {}^t\mathrm{Var}^{-1}$, $I = -{}^t\mathrm{Var}^{-1} - \mathrm{Var}^{-1}$, $M = -\,\mathrm{Var}\,{}^t\mathrm{Var}^{-1}$.
- Gabriélov polar method and Escario's give I for a DVBC. A'Campo can work with integral bilinear forms of high rank.

A'Campo's question

Is ρ_* conjugated to its inverse (over $\mathbb{Z}$)?

- The answer is YES over $\mathbb{Q}$ (Monodromy theorem).



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.
- Var upper triangular with diagonals 1, Vart = ^{t}Var.
- $S = {}^t\mathrm{Var}^{-1}$, $I = -{}^t\mathrm{Var}^{-1} - \mathrm{Var}^{-1}$, $M = -\,\mathrm{Var}\,{}^t\mathrm{Var}^{-1}$.
- Gabriélov polar method and Escario's give I for a DVBC. A'Campo can work with integral bilinear forms of high rank.

A'Campo's question

Is ρ_* conjugated to its inverse (over $\mathbb{Z}$)?

- The answer is YES over $\mathbb{Q}$ (Monodromy theorem).
- The answer is YES for curves: any topological type can be realized over the reals and take complex conjugation.



Picard-Lefschetz

- From a morsification $(\Delta_1, \ldots, \Delta_\mu)$ *distinguished basis of vanishing cycles* (DBVC) of $H_2(F;\mathbb{Z})$,

$$\Delta_i \cdot \Delta_i = -2.$$

  $(\nabla_1, \ldots, \nabla_\mu)$ dual basis of $H_2(F, \partial F;\mathbb{Z})$
- Matrices of var, ${}^t\mathrm{var}, L, \cdot, \rho_*$: Var, Vart, S, I, M.
- Var upper triangular with diagonals 1, Vart = ^{t}Var.
- $S = {}^t\mathrm{Var}^{-1}$, $I = -{}^t\mathrm{Var}^{-1} - \mathrm{Var}^{-1}$, $M = -\,\mathrm{Var}\,{}^t\mathrm{Var}^{-1}$.
- Gabriélov polar method and Escario's give I for a DVBC. A'Campo can work with integral bilinear forms of high rank.

A'Campo's question

Is ρ_* conjugated to its inverse (over $\mathbb{Z}$)?

- The answer is YES over $\mathbb{Q}$ (Monodromy theorem).
- The answer is YES for curves: any topological type can be realized over the reals and take complex conjugation.
- Possible counterexample: look for a SIS such that the topological type of the tangent cone cannot be realized over the reals.



Surfaces in $\mathbb{C}^3$

Superisolated and Lê-Yomdin singularities

Zariski tuples and Jordan form

Zariski tuples of non-Alexander type

Quasi-homogeneous and Weighted Lê-Yomdin singularities



Weighted Lê-Yomdin singularities

Definition

$\omega \in \mathbb{Z}^3_{>0}$ coprime, $(S,0) = \{F = 0\}$, $F \in \mathbb{C}\{x,y,z\}$, isolated surface singularity. Decompose F in ω-quasi-homogeneous forms:

$$F(x,y,z) = \overbrace{F_m(x,y,z)}^{\neq 0} + F_{m+k}(x,y,z) + \dots$$

Let $C_j := V_{\mathbb{P}}(F_m) \subset \mathbb{P}^2_\omega$ (as orbifold). Then S is a (k,ω)-*weighted Lê-Yomdin singularity* (WLYS) if at each $P \in C_m \cap C_{m+k}$ the *local equation* of C_m at P is of order 1.



Weighted Lê-Yomdin singularities

Definition

$\omega \in \mathbb{Z}^3_{>0}$ coprime, $(S,0) = \{F = 0\}$, $F \in \mathbb{C}\{x,y,z\}$, isolated surface singularity. Decompose F in ω-quasi-homogeneous forms:

$$F(x,y,z) = \overbrace{F_m(x,y,z)}^{\neq 0} + F_{m+k}(x,y,z) + \dots$$

Let $C_j := V_{\mathbb{P}}(F_m) \subset \mathbb{P}^2_\omega$ (as orbifold). Then S is a (k,ω)-*weighted Lê-Yomdin singularity* (WLYS) if at each $P \in C_m \cap C_{m+k}$ the *local equation* of C_m at P is of order 1.

Weighted blow-ups



Weighted Lê-Yomdin singularities

Definition

$\omega \in \mathbb{Z}^3_{>0}$ coprime, $(S,0) = \{F = 0\}$, $F \in \mathbb{C}\{x,y,z\}$, isolated surface singularity. Decompose F in ω-quasi-homogeneous forms:

$$F(x,y,z) = \overbrace{F_m(x,y,z)}^{\neq 0} + F_{m+k}(x,y,z) + \dots$$

Let $C_j := V_{\mathbb{P}}(F_m) \subset \mathbb{P}^2_\omega$ (as orbifold). Then S is a (k,ω)-*weighted Lê-Yomdin singularity* (WLYS) if at each $P \in C_m \cap C_{m+k}$ the *local equation* of C_m at P is of order 1.

Weighted blow-ups

- Rephrase Lê-Yomdin singularity work replacing the first blow-up by an ω-weighted blow-up.



Weighted Lê-Yomdin singularities

Definition

$\omega \in \mathbb{Z}^3_{>0}$ coprime, $(S,0) = \{F = 0\}$, $F \in \mathbb{C}\{x,y,z\}$, isolated surface singularity. Decompose F in ω-quasi-homogeneous forms:

$$F(x,y,z) = \overbrace{F_m(x,y,z)}^{\neq 0} + F_{m+k}(x,y,z) + \dots$$

Let $C_j := V_{\mathbb{P}}(F_m) \subset \mathbb{P}^2_\omega$ (as orbifold). Then S is a (k,ω)-*weighted Lê-Yomdin singularity* (WLYS) if at each $P \in C_m \cap C_{m+k}$ the *local equation* of C_m at P is of order 1.


Weighted blow-ups

- Rephrase Lê-Yomdin singularity work replacing the first blow-up by an ω-weighted blow-up.
- Notion of Zariski tuples can be extended.



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle

- $f_t = z^{12} + xy^3z + tx^3y^2 + x^6 = 0$ equisingular deformation.



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle

- $f_t = z^{12} + xy^3z + tx^3y^2 + x^6 = 0$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{[0:1:0]_\omega\} \subset \mathbb{P}^2_\omega$, intersection number with $X = 0$ changes.



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle

- $f_t = z^{12} + xy^3z + tx^3y^2 + x^6 = 0$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{[0:1:0]_\omega\} \subset \mathbb{P}^2_\omega$, intersection number with $X = 0$ changes.
- $F_t = f_t + y^5 = 0$ 3-WLYS, same topology and distinct tangent cone.



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle

- $f_t = z^{12} + xy^3z + tx^3y^2 + x^6 = 0$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{[0:1:0]_\omega\} \subset \mathbb{P}^2_\omega$, intersection number with $X = 0$ changes.
- $F_t = f_t + y^5 = 0$ 3-WLYS, same topology and distinct tangent cone.

Non-linear deformations



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle

- $f_t = z^{12} + xy^3z + tx^3y^2 + x^6 = 0$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{[0:1:0]_\omega\} \subset \mathbb{P}^2_\omega$, intersection number with $X = 0$ changes.
- $F_t = f_t + y^5 = 0$ 3-WLYS, same topology and distinct tangent cone.

Non-linear deformations

- $f_t = (y^2 + x^3)^2 + (yz + tx^2)^3$ equisingular deformation.



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS

- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle

- $f_t = z^{12} + xy^3z + tx^3y^2 + x^6 = 0$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{[0:1:0]_\omega\} \subset \mathbb{P}^2_\omega$, intersection number with $X = 0$ changes.
- $F_t = f_t + y^5 = 0$ 3-WLYS, same topology and distinct tangent cone.

Non-linear deformations

- $f_t = (y^2 + x^3)^2 + (yz + tx^2)^3$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{P = [0:0:1], Q_t = [-1:1:-t]\}$ (ordinary cusp at Q_t)



Equisingular families in $\mathbb{P}^2_\omega$, $\omega = (2,3,1)$ and WLYS

Briançon-Speder WYLS


- $F_t = y^5 + tx^6y + zx^7 + z^{15} = 0$, equisingular deformation of smooth curves $\Rightarrow$ WYLS
- Tangent line at $P := [1:0:0]_\omega$ is $ty + z = 0$.

Tangent cone, E. A., J. FdB, I. Luengo, A. Melle

- $f_t = z^{12} + xy^3z + tx^3y^2 + x^6 = 0$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{[0:1:0]_\omega\} \subset \mathbb{P}^2_\omega$, intersection number with $X = 0$ changes.
- $F_t = f_t + y^5 = 0$ 3-WLYS, same topology and distinct tangent cone.

Non-linear deformations

- $f_t = (y^2 + x^3)^2 + (yz + tx^2)^3$ equisingular deformation.
- $\mathrm{Sing}\{f_t = 0\} = \{P = [0:0:1], Q_t = [-1:1:-t]\}$ (ordinary cusp at Q_t)
- $F_t = f_t + z^{13} + x^2y^3$ 1-WLYS same topology.



Thanks for your attention!

